

Quantum Spheres As Groupoid C^* -algebras

Albert J. L. Sheu*
 Department of Mathematics
 University of Kansas
 Lawrence, KS 66045-2142
 U. S. A.

Abstract

In this paper, we show that the C^* -algebras of quantum spheres can be realized as concrete groupoid C^* -algebras.

1 Introduction

Ever since the successful use of groupoid C^* -algebras [Re] by Connes in studying the interaction between operator algebras and foliation geometry [Co], it has been well recognized that the theory of groupoid C^* -algebras provides a very useful tool and a nearly universal context for the study of operator algebras, especially in connection with geometric structures. The results obtained in [CuM, MRe, SaShU, Sh1, Sh4] are a few examples supporting this aspect. In particular, it has been found that some interesting C^* -algebras associated with singular foliations can be effectively described and studied in the context of C^* -algebras of (essentially) discrete groupoids [Sh3, Sh4].

Recent fast developments in deformation quantization [Ri1, Ri2, Sh2] including quantum groups [Dr, RTF, So, Wo1, Wo2, Ri3] showed that the problem of quantization of Poisson structures and group structures is intimately related to the underlying singular symplectic foliation [We, LuWe]. Furthermore in the paper [Sh4], it is found that the C^* -algebra arising from deformation quantization using quantum groups can be studied by the method of groupoids.

In fact, it is shown [Sh4] that C^* -algebras $C(SU(n)_q)$ of the quantum groups $SU(n)_q$ can be realized as a C^* -subalgebra of some groupoid C^* -algebras. We also know that $C(SU(2)_q) \simeq C^*(\mathfrak{F})$ where

$$\mathfrak{F} = \{(m, j, k) \mid \text{if } k = \infty, \text{ then } m = j\}$$

is a subgroupoid of the Toeplitz groupoid $\mathbb{Z} \times (\mathbb{Z} \times \overline{\mathbb{Z}}|_{\overline{\mathbb{Z}}_{\geq}})$.

In the study of quantizing Poisson Lie groups, it is also of great interest to see how the Poisson structures reduce to homogeneous spaces and how the

*Partially supported by NSF-Grant DMS-9303231

resulting Poisson spaces can be quantized [DoGu]. Fundamental examples are provided by the study of quantum spheres [Po, VSo]. It is also found that C*-algebras arising from such a quantization are closely related to the underlying symplectic structure and the corresponding groupoid [Sh4]. In particular, we know that $C(S_q^{2n+1})$ of the quantum sphere S_q^{2n+1} [VSo] can be realized as a C*-subalgebra of some groupoid C*-algebra, and $C(S_{\mu c}^2) \simeq C^*(\mathfrak{G})$ where $S_{\mu c}^2$ is Podles' quantum sphere [Po] with $c > 0$ and

$$\mathfrak{G} = \{(j, j, k_1, k_2) \mid k_1 = \infty \text{ or } k_2 = \infty\}$$

is a subgroupoid of the 2-dimensional Toeplitz groupoid $\mathbb{Z}^2 \times \overline{\mathbb{Z}}^2|_{\overline{\mathbb{Z}}^2_{\geq}}$.

However it is not clear and it is an interesting question whether other $C(S_q^{2n+1})$ or $C(SU(n)_q)$ can be realized as a groupoid C*-algebra itself instead of a C*-subalgebra of some groupoid C*-algebra. In this paper, we give a positive answer to this question for $C(S_q^{2n+1})$ and show that the underlying groupoid can be identified explicitly and is independent of the parameter q . This also gives a proof of the fact that for a fixed n , the C*-algebras $C(S_q^{2n+1})$ are isomorphic and independent of q . We mention that the above statements are also true for quantum $SU(3)$, but the proof is more complicated and we will deal with it in a separate paper.

2 Quantum sphere and groupoid

We refer to the paper [Sh4] for notations and concepts.

Recall that the C*-algebra of quantum group $SU(n)_q$ is generated by elements u_{ij} satisfying certain commutation relations and the C*-algebra of quantum spheres $S_q^{2n+1} = SU(n)_q \setminus SU(n+1)_q$ defined as homogeneous quantum spaces [N] can be identified with

$$C(S_q^{2n+1}) = C^*(\{u_{n+1,m} \mid 1 \leq m \leq n+1\}).$$

In [Sh4], we give explicit embeddings of $C(SU(n)_q)$ and $C(S_q^{2n+1})$ into concrete groupoid C*-algebras.

Recall that $C(S_q^{2n+1})$ can be embedded in the groupoid C*-algebra $C^*(\mathcal{F}^n)$, where $\mathcal{F}^n$ is the n -dimensional Toeplitz groupoid $\mathbb{Z} \times (\mathbb{Z}^n \times \overline{\mathbb{Z}}^n|_{\overline{\mathbb{Z}}^n_{\geq}})$. Let

$$F_i := \{w \in F : w_i = \infty\} \cong \overline{\mathbb{Z}}_{\geq}^{n-1}$$

denote a 'face' of the unit space $F \cong \overline{\mathbb{Z}}_{\geq}^n$ of the groupoid $\mathcal{F}^n$ and $\partial F = \cup_{i=1}^n F_i$ be the 'boundary' of F .

Let $\rho_{i*} : \mathcal{F}^n|_{F_i} \rightarrow \mathcal{F}^n$ be the proper continuous groupoid inclusion map and

$$\rho_i = C^*(\rho_{i*}) : C^*(\mathcal{F}^n) \rightarrow C^*(\mathcal{F}^n)|_{F_i} = C^*(\mathcal{F}^n|_{F_i})$$

be the corresponding C*-algebra homomorphism implementing the restriction to the face F_i . Similarly we consider the embeddings $\rho_{\partial F*} : \mathcal{F}^n|_{\partial F} \rightarrow \mathcal{F}^n$ and

$\rho_{i,\partial F*} : \mathcal{F}^n|_{F_i} \rightarrow \mathcal{F}^n|_{\partial F}$, and the corresponding algebra homomorphisms $\rho_{\partial F}$ and $\rho_{i,\partial F}$.

Now the restriction of $C(S_q^{2n+1})$ to the face F_n dominates the restriction to any other F_i , in the sense that ρ_i on $C(S_q^{2n+1})$ factors through ρ_n on $C(S_q^{2n+1})$. More precisely, we have the following lemma.

Lemma 1 $\rho_{n,\partial F}$ when restricted to $C(S_q^{2n+1})|_{\partial F}$ is a C^* -algebra isomorphism

$$\rho_{n,\partial F} = C^*(\rho_{n,\partial F*}) : C(S_q^{2n+1})|_{\partial F} \rightarrow C(S_q^{2n+1})|_{F_n}$$

with its inverse

$$\pi_{n,\partial F} = C^*(\pi_{n,\partial F*}) : C(S_q^{2n+1})|_{F_n} \rightarrow C(S_q^{2n+1})|_{\partial F}$$

arising from the proper continuous groupoid quotient map $\pi_{n,\partial F*} : \mathcal{F}^n|_{\partial F} \rightarrow \mathcal{F}^n|_{F_n}$ sending (z, x, w) to (z, x, w') where $w'_n = \infty$ and $w'_j = w_j$ for all $j < n$.

Proof.

In fact, since the restriction $\mathcal{A}|_{\partial F}$ of any C^* -subalgebra $\mathcal{A} \subset C^*(\mathcal{F}^n)$ can be identified, under the map $\oplus_{i=1}^n \rho_{i,\partial F} : \mathcal{A}|_{\partial F} \rightarrow \oplus_{i=1}^n \mathcal{A}|_{F_i}$, with the pull-back

$$\mathcal{A}' = \{\oplus_i a_i \in \oplus_i \mathcal{A}|_{F_i} : \rho_{ij}(a_j) = \rho_{ji}(a_i) \text{ for all } i, j\}$$

of the homomorphisms $\mathcal{A}|_{F_j} \xrightarrow{\rho_{ij}} \mathcal{A}|_{F_i \cap F_j}$ arising from the proper continuous embedding ρ_{ij*} of the face $F_i \cap F_j$ into the face F_i , we only need to show that in the commuting diagram

$$\begin{array}{ccc} \mathcal{A}|_{\partial F} & \xrightarrow{\rho_{i,\partial F}} & \mathcal{A}|_{F_i} \\ \rho_{n,\partial F} \downarrow & & \downarrow \rho_{ni} \\ \mathcal{A}|_{F_n} & \xrightarrow{\rho_{in}} & \mathcal{A}|_{F_i \cap F_n} \end{array}$$

with $\mathcal{A} = C(S_q^{2n+1})$, ρ_{ni} is an isomorphism for $i < n$ with the inverse

$$\pi_{ni} = C^*(\pi_{ni*}) : C(S_q^{2n+1})|_{F_i \cap F_n} \rightarrow C(S_q^{2n+1})|_{F_i}$$

arising from the groupoid quotient map $\pi_{ni*} : \mathcal{F}^n|_{F_i} \rightarrow \mathcal{F}^n|_{F_i \cap F_n}$ sending (z, x, w) to (z, x, w') where $w'_j = w_j$ for all $j < n$ and $w'_n = \infty$. Note that $\pi_{ni} \circ \rho_{in} = C^*(\pi_{n,\partial F*}|_{F_i})$.

It is clear that $\rho_{ni} \circ \pi_{ni} = Id$ since $\pi_{ni*} \circ \rho_{ni*} = Id$. On the other hand, we can verify $\pi_{ni} \circ \rho_{ni} = Id$ on the generators $\rho_i(u_{n+1,m})$ of $C(S_q^{2n+1})|_{F_i}$. Recall that $u_{n+1,m} \in C(S_q^{2n+1})$ is identified with

$$(\tau_{n+1} \otimes \pi_{L_1})(\Delta u_{n+1,m}) = t_{n+1} \otimes \underbrace{\gamma \otimes \dots \otimes \gamma}_{n+1-m} \otimes \underbrace{\alpha^* \otimes 1 \otimes \dots \otimes 1}_{m-1}$$

in $C^*(\mathcal{F}^n)$. It is easy to see that $\rho_i((\tau_{n+1} \otimes \pi_{L_1})\Delta(u_{n+1,m})) = 0$ for $m \leq n+1-i$, and hence we only need to show that $(\pi_{ni} \circ \rho_{ni})(\rho_i(u_{n+1,m})) = \rho_i(u_{n+1,m})$ for $m > n+1-i$. But

$$\rho_i((\tau_{n+1} \otimes \pi_{L_1})(\Delta u_{n+1,n+2-i})) = t_{n+1} \otimes \underbrace{\gamma \otimes \dots \otimes \gamma}_{i-1} \otimes \underbrace{s_i^{-1} \otimes 1 \otimes \dots \otimes 1}_{m-1}$$

with s_i the canonical generator of $C(\mathbb{T}) \cong C^*(\{\infty\} \times \mathbb{Z})$, and

$$\rho_i((\tau_{n+1} \otimes \pi_{L_1})(\Delta u_{n+1,m})) = t_{n+1} \otimes \underbrace{\gamma \otimes \dots \otimes \gamma}_{n+1-m} \otimes \underbrace{\alpha^* \otimes 1 \otimes \dots \otimes 1}_{m-1}$$

for $m > n + 2 - i$, are of the form $a \otimes 1$ where $a \in C^*(\mathcal{F}^{n-1}|_{F_i})$ and $1 \in C^*(\mathbb{Z} \times \overline{\mathbb{Z}}|_{\overline{\mathbb{Z}}_{\geq}})$. Since clearly $(\pi_{ni} \circ \rho_{ni})(a \otimes 1) = a \otimes 1$, we get $(\pi_{ni} \circ \rho_{ni})(\rho_i(u_{n+1,m})) = \rho_i(u_{n+1,m})$ for $m > n + 1 - i$, and hence $Id = \pi_{ni} \circ \rho_{ni}$ on $\rho_i(C(S_q^{2n+1})) = C(S_q^{2n+1})|_{F_i}$ for $i < n$. (Note that π_{ni} and $\pi_{n,\partial F}$ are well-defined because $C(S_q^{2n+1})|_{F_i} \subset C^*(\mathcal{F}^{n-1}|_{F_i}) \otimes 1$ for $i < n$.)

■

3 The groupoid realization of quantum sphere

By the fact [Sh4] that $C(\mathbb{T}) \otimes \mathcal{K}(\ell^2(\mathbb{Z}_{\geq}^n)) \cong C^*(\mathcal{F}^n|_{\mathbb{Z}_{\geq}^n}) \subset C(S_q^{2n+1})$, we have the commuting diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & C(\mathbb{T}) \otimes \mathcal{K}(\ell^2(\mathbb{Z}_{\geq}^n)) & \rightarrow & C(S_q^{2n+1}) & \xrightarrow{\rho_{\partial F}} & C(S_q^{2n+1})|_{\partial F} \rightarrow 0 \\ & & \parallel & & \cap & & \cap \\ 0 & \rightarrow & C(\mathbb{T}) \otimes \mathcal{K}(\ell^2(\mathbb{Z}_{\geq}^n)) & \rightarrow & C^*(\mathcal{F}^n) & \xrightarrow{\rho_{\partial F}} & C^*(\mathcal{F}^n|_{\partial F}) \rightarrow 0 \end{array}$$

with exact rows. It is clear that the structure of $C(S_q^{2n+1})$ is completely determined by $C(S_q^{2n+1})|_{\partial F} \subset C^*(\mathcal{F}^n|_{\partial F})$. On the other hand, $C(S_q^{2n+1})|_{\partial F} = \pi_{n,\partial F}(C(S_q^{2n+1})|_{F_n})$ can be determined explicitly when $C(S_q^{2n+1})|_{F_n}$ is.

Define a subquotient groupoid $\mathfrak{F}_n$ of $\mathcal{F}^n = \mathbb{Z} \times (\mathbb{Z}^n \times \overline{\mathbb{Z}}^n|_{\overline{\mathbb{Z}}_{\geq}^n})$ as follows. Let

$$\widetilde{\mathfrak{F}}_n := \{(z, x, w) \in \mathcal{F}^n \mid w_i = \infty \implies$$

$$x_i = -z - x_1 - x_2 - \dots - x_{i-1} \text{ and } x_{i+1} = \dots = x_n = 0\}$$

be a subgroupoid of $\mathcal{F}^n$. Define $\mathfrak{F}_n := \widetilde{\mathfrak{F}}_n / \sim$ where $\sim$ is the equivalence relation generated by

$$(z, x, w) \sim (z, x, w_1, \dots, w_i = \infty, \infty, \dots, \infty)$$

for all (z, x, w) with $w_i = \infty$ for an $1 \leq i \leq n$.

Theorem 2 $C(S_q^{2n+1}) \simeq C^*(\mathfrak{F}_n)$.

Proof.

Actually we prove by induction on n that

$$C(S_q^{2n+1}) = \iota_n \eta_n(C^*(\mathfrak{F}_n)) \simeq C^*(\mathfrak{F}_n)$$

where η_n is the injective map arising from the groupoid quotient map $\eta_{n*} : \widetilde{\mathfrak{F}}_n \rightarrow \mathfrak{F}_n$ and $\iota_n : C^*(\widetilde{\mathfrak{F}}_n) \rightarrow C^*(\mathcal{F}^n)$ implements the inclusion of the open subgroupoid $\widetilde{\mathfrak{F}}_n$ in $\mathcal{F}^n$.

When $n = 1$, the equivalence relation $\sim$ is trivial, i.e. $\widetilde{\mathfrak{F}}_1 = \widetilde{\mathfrak{F}}_1 / \sim = \mathfrak{F}_1$ and hence $\iota_1 \eta_1 (C^*(\mathfrak{F}_1)) = \iota_1 (C^*(\widetilde{\mathfrak{F}}_1))$ with

$$\widetilde{\mathfrak{F}}_1 = \{(m, j, k) \mid \text{if } k = \infty, \text{ then } m = j\}.$$

On the other hand, the C*-algebra $C(S_q^{2n+1}) = C(SU(2)_q) \subset C^*(\mathcal{F}^1)$ is generated by

$$(\tau_2 \otimes \pi_{L_1})(u_{2,1}) = t_2 \otimes \gamma$$

and

$$(\tau_2 \otimes \pi_{L_1})(u_{2,2}) = t_2 \otimes \alpha^*,$$

and hence coincides with $\iota_1 (C^*(\widetilde{\mathfrak{F}}_1))$ as can be easily verified.

Now we assume the induction hypothesis

$$C(S_q^{2n-1}) = \iota_{n-1} \eta_{n-1} (C^*(\mathfrak{F}_{n-1})) \simeq C^*(\mathfrak{F}_{n-1}).$$

By identifying $(z, x, w) \in \mathcal{F}^{n-1} \subset \mathbb{Z} \times \mathbb{Z}^{n-1} \times \overline{\mathbb{Z}}_{\geq}^{n-1}$ with $(z, x, 0, w, \infty) \in \mathcal{F}^n|_{F_n} \subset \mathbb{Z} \times \mathbb{Z}^n \times \overline{\mathbb{Z}}_{\geq}^{n-1} \times \{\infty\}$, we regard $\mathcal{F}^{n-1}$ and hence $\widetilde{\mathfrak{F}}_{n-1}$ as open subgroupoids of $\mathcal{F}^n|_{F_n}$, and regard $C^*(\mathcal{F}^{n-1})$ and $C^*(\widetilde{\mathfrak{F}}_{n-1})$ as C*-subalgebras of $C^*(\mathcal{F}^n)$. Note that under this identification,

$$\widetilde{\mathfrak{F}}_{n-1} := \{(z, x, w) \in \mathcal{F}^n \mid w_n = \infty, x_n = 0, \text{ and for } i < n,$$

$$\text{"}w_i = \infty \implies x_i = -z - x_1 - \dots - x_{i-1} \text{ and } x_{i+1} = \dots = x_n = 0\text{"}\}$$

and $\mathfrak{F}_{n-1} = \widetilde{\mathfrak{F}}_{n-1} / \sim$ with respect to the same equivalence relation $\sim$.

Now note that $C(S_q^{2n+1})|_{F_n} \cong C(S_q^{2n-1}) \subset C^*(\mathcal{F}^{n-1})$ under the automorphism ϕ on $C^*(\mathcal{F}^n|_{F_n}) \supset C^*(\mathcal{F}^{n-1})$, corresponding to the groupoid isomorphism ϕ_* on $\mathcal{F}^n|_{F_n} \subset \mathbb{Z} \times \mathbb{Z}^n \times \overline{\mathbb{Z}}_{\geq}^{n-1}$ defined by

$$\phi_*(z, x, w) = \left(z, x + \left(0, \dots, 0, -z - \sum_{i=1}^{n-1} x_i \right), w \right).$$

More precisely, $C(S_q^{2n+1})|_{F_n} = \phi (C(S_q^{2n-1}))$.

Thus from the induction hypothesis, we get

$$C(S_q^{2n+1})|_{F_n} = \phi [\iota_{n-1} \eta_{n-1} (C^*(\mathfrak{F}_{n-1}))].$$

Since under the groupoid isomorphism ϕ_* , $\widetilde{\mathfrak{F}_{n-1}}$ is mapped to the open subgroupoid

$$\widetilde{\mathfrak{F}'_{n-1}} := \phi_* \left(\widetilde{\mathfrak{F}_{n-1}} \right) = \{(z, x, w) \in \mathcal{F}^n \mid w_n = \infty, x_n = -z - \sum_{i=1}^{n-1} x_i,$$

and for $i < n$, “ $w_i = \infty \implies x_i = -z - x_1 - \dots - x_{i-1}$ and $x_{i+1} = \dots = x_n = 0$ ”}

$$= \{(z, x, w) \in \mathcal{F}^n \mid w_n = \infty, \text{ and for any } i \leq n,$$

$$\text{“}w_i = \infty \implies x_i = -z - x_1 - \dots - x_{i-1} \text{ and } x_{i+1} = \dots = x_n = 0\text{”}\}$$

of $\mathcal{F}^n|_{F_n}$ and the equivalence relation $\sim$ is preserved, it is clear that

$$C(S_q^{2n+1})|_{F_n} = \iota'_{n-1} \eta'_{n-1} (C^*(\mathfrak{F}'_{n-1}))$$

where $\eta'_{n-1*} : \widetilde{\mathfrak{F}'_{n-1}} \rightarrow \mathfrak{F}'_{n-1} := \widetilde{\mathfrak{F}'_{n-1}} / \sim$ is the quotient map of the subgroupoid $\widetilde{\mathfrak{F}'_{n-1}}$ of $\mathcal{F}^n|_{F_n}$, and $\iota'_{n-1} : C^*(\widetilde{\mathfrak{F}'_{n-1}}) \rightarrow C^*(\mathcal{F}^n|_{F_n})$ is the inclusion.

Now for $(z, x, w) \in \widetilde{\mathfrak{F}'_{n-1}}$, we have $w_n = \infty$, and if $w_i = \infty$ for some $i < n$ then $x_n = 0$. Recall that $\pi_{n, \partial F*}(z, x, w) = (z, x, w')$ with $w' = (w_1, \dots, w_{n-1}, \infty)$. We claim that

$$\widetilde{\mathfrak{F}''_{n-1}} := \pi_{n, \partial F*}^{-1} \left(\widetilde{\mathfrak{F}'_{n-1}} \right) = \{(z, x, w) \in \mathcal{F}^n|_{\partial F} : \text{for any } i \leq n, \}$$

$$\text{“}w_i = \infty \implies x_i = -z - x_1 - \dots - x_{i-1} \text{ and } x_{i+1} = \dots = x_n = 0\text{”}\}.$$

Indeed, if $(z, x, w) \in \mathcal{F}^n|_{\partial F}$ and $\pi_{n, \partial F*}(z, x, w) \in \widetilde{\mathfrak{F}'_{n-1}}$, then (z, x, w) satisfies the above condition since it is already satisfied by $(z, x, w') \in \widetilde{\mathfrak{F}'_{n-1}}$ and w does not have more ∞ entries than w' . On the other hand, if $(z, x, w) \in \mathcal{F}^n|_{\partial F}$ satisfies the above condition, then either $(z, x, w) \in \mathcal{F}^n|_{F_n}$ and hence $(z, x, w') = (z, x, w)$ since $w_n = \infty$, or $w_i = \infty$ for some $i < n$ which implies that $x_i = -z - x_1 - \dots - x_{i-1}$ with $x_{i+1} = \dots = x_n = 0$ and hence

$$x_n = 0 = (-z - x_1 - \dots - x_{i-1}) - x_i = -z - x_1 - \dots - x_{n-1}.$$

In either case, the above condition is satisfied by $(z, x, w') = \pi_{n, \partial F*}(z, x, w) \in \mathcal{F}^n|_{F_n}$, and hence $\pi_{n, \partial F*}(z, x, w) \in \widetilde{\mathfrak{F}'_{n-1}}$.

With $\widetilde{\mathfrak{F}'_{n-1}}$ canonically embedded in $\mathcal{F}^n|_{F_n}$, the composition

$$\eta''_{n-1*} := \eta'_{n-1*} \circ \pi_{n, \partial F*} : \widetilde{\mathfrak{F}''_{n-1}} \rightarrow \mathfrak{F}'_{n-1}$$

of quotient maps realizes $\mathfrak{F}'_{n-1}$ as a quotient groupoid $\widetilde{\mathfrak{F}''_{n-1}}/\sim$ of $\widetilde{\mathfrak{F}''_{n-1}}$ where the equivalence relation $\sim$ is the original one extended to $\widetilde{\mathfrak{F}''_{n-1}}$, i.e. the relation generated by

$$(z, x, w) \sim (z, x, w_1, \dots, w_i = \infty, \infty, \dots, \infty)$$

for all (z, x, w) with $w_i = \infty$ for an $1 \leq i \leq n$. Since $\pi_{n, \partial F} : \widetilde{\mathfrak{F}''_{n-1}} \rightarrow \widetilde{\mathfrak{F}'_{n-1}} \subset \mathcal{F}^n|_{F_n}$ where the inclusion is implemented by $\iota'_{n-1} : C^*(\widetilde{\mathfrak{F}'_{n-1}}) \rightarrow C^*(\mathcal{F}^n|_{F_n})$ at the C*-algebra level, it is easy to see that $\eta''_{n-1} = \pi_{n, \partial F} \circ \eta'_{n-1}$ and hence

$$\begin{aligned} C(S_q^{2n+1})|_{\partial F} &= \pi_{n, \partial F} (C(S_q^{2n+1})|_{F_n}) \\ &= \pi_{n, \partial F} (\eta'_{n-1} (C^*(\mathfrak{F}'_{n-1}))) = \eta''_{n-1} (C^*(\mathfrak{F}'_{n-1})) \subset C^*(\mathcal{F}^n|_{\partial F}). \end{aligned}$$

Note that $\widetilde{\mathfrak{F}''_{n-1}} \subset \mathcal{F}^n|_{\partial F}$ is a closed subgroupoid of $\widetilde{\mathfrak{F}_n} \subset \mathcal{F}^n$ and hence $\mathfrak{F}'_{n-1}$ is a closed subgroupoid of $\mathfrak{F}_n$ with η''_{n-1} equal to the restriction of η_n to $\widetilde{\mathfrak{F}''_{n-1}}$. On the other hand, on the ‘interior’ $\widetilde{\mathfrak{F}_n} - \widetilde{\mathfrak{F}''_{n-1}} = \mathcal{F}^n - (\mathcal{F}^n|_{\partial F})$, the relation $\sim$ is trivial and hence

$$\begin{array}{ccccccc} & & & & C(S_q^{2n+1})|_{\partial F} & & \\ & & & & \parallel & & \\ 0 & \rightarrow & \iota_n \eta_n (C^*(\mathfrak{F}_n - \mathfrak{F}'_{n-1})) & \subset & \iota_n \eta_n (C^*(\mathfrak{F}_n)) & \xrightarrow{\rho_{\partial F}} & \eta''_{n-1} (C^*(\mathfrak{F}'_{n-1})) \rightarrow 0 \\ & & \parallel & & \cap & & \cap \\ 0 & \rightarrow & \iota_n C^*(\widetilde{\mathfrak{F}_n} - \widetilde{\mathfrak{F}''_{n-1}}) & \subset & \iota_n C^*(\widetilde{\mathfrak{F}_n}) & \xrightarrow{\rho_{\partial F}} & C^*(\widetilde{\mathfrak{F}''_{n-1}}) \rightarrow 0 \\ & & \parallel & & \cap & & \cap \\ 0 & \rightarrow & C(\mathbb{T}) \otimes \mathcal{K}(\ell^2(\mathbb{Z}_{\geq}^n)) & \subset & C^*(\mathcal{F}^n) & \xrightarrow{\rho_{\partial F}} & C^*(\mathcal{F}^n|_{\partial F}) \rightarrow 0 \end{array}$$

But as we have seen $C(S_q^{2n+1})$ is the only C*-subalgebra of $C^*(\mathcal{F}^n)$ whose image under $\rho_{\partial F}$ coincides with $C(S_q^{2n+1})|_{\partial F} \subset C^*(\mathcal{F}^n|_{\partial F})$. So we get $\iota_n \eta_n (C^*(\mathfrak{F}_n)) = C(S_q^{2n+1}) \subset C^*(\mathcal{F}^n)$ as stated. $\blacksquare$

Corollary 3 *The C*-algebra $C(S_q^{2n+1})$ is independent of q .*

We remark that this fact can also be derived from Vaksman and Soibelman’s result [VSo].

4 Bibliography

References

- [Co] A. Connes, *A survey of foliation and operator algebras*, Proc. Symp. Pure Math. Vol. 38, Part I, AMS, Providence, 1982, 521-628.

- [CuM] R. E. Curto and P. S. Muhly, *C*-algebras of multiplication operators on Bergman spaces*, J. Func. Anal. 64 (1985), 315-329.
- [DoGu] J. Donin and D. Gurevich, *Quantum orbits of the R-matrix type*, Lett. Math. Phys. 35 (1995), 263-276.
- [Dr] V. G. Drinfeld, *Quantum groups*, Proc. I.C.M. Berkeley 1986, Vol. 1, 789-820, Amer. Math. Soc., Providence, 1987.
- [LuWe] J. H. Lu and A. Weinstein, *Poisson Lie groups, dressing transformations and Bruhat decompositions*, J. Diff. Geom. 31 (1990), 501-526.
- [MRe] P. S. Muhly and J. N. Renault, *C*-algebras of multivariable Wiener-Hopf operators*, Trans. Amer. Math. Soc. 274 (1982), 1-44.
- [N] G. Nagy, *On the Haar measure of the quantum $SU(N)$ group*, Comm. Math. Phys. 153 (1993), 217-228.
- [Po] P. Podles, *Quantum spheres*, Letters Math. Phys. 14 (1987), 193-202.
- [Re] J. Renault, *A Groupoid Approach to C*-algebras*, Lecture Notes in Mathematics, Vol. 793, Springer-Verlag, New York, 1980.
- [RTF] N. Yu. Reshetikhin, L. A. Takhtadzhyan, and L. D. Faddeev, *Quantization of Lie groups and Lie algebras*, Leningrad Math. J. 1 (1990), 193-225.
- [Ri1] M. A. Rieffel, *Deformation quantization and operator algebras*, in "Proc. Symp. Pure Math., Vol. 51", AMS, Providence, 1990, pp. 411-423.
- [Ri2] ———, *Deformation quantization for actions of $\mathbb{R}^d$* , Memoirs of AMS, Vol. 106, No. 506, 1993.
- [Ri3] ———, *Compact quantum groups associated with toral subgroups*, in "Contemporary Mathematics", Vol. 145, AMS, Providence, 1993, pp. 465-491.
- [SaShU] N. Salinas, A. J. L. Sheu, and H. Upmeyer, *Toeplitz operators on pseudoconvex domains and foliation C*-algebras*, Annals of Mathematics, 130 (1989), 531-565.
- [Sh1] A. J. L. Sheu, *Reinhardt domains, boundary geometry and Toeplitz C*-algebras*, Journal of Functional Analysis, 92 (1990), 264-311.
- [Sh2] ———, *Quantization of the Poisson $SU(2)$ and its Poisson homogeneous space – the 2-sphere*, Comm. Math. Phys. 135 (1991), 217-232.
- [Sh3] ———, *Some examples of singular foliation C*-algebras*, in "Current Topics in Operator Algebras", H. Araki et al (Ed.s), World Scientific, Singapore, 1991, pp. 326-340.
- [Sh4] ———, *Compact quantum groups and groupoid C*-algebras*, J. Func. Anal. 144 (1997), 371-393.

- [So] Ya. S. Soibelman, *The algebra of functions on a compact quantum group, and its representations*, Algebra Analiz. 2 (1990), 190-221. (Leningrad Math. J., 2 (1991), 161-178.)
- [VSo] L. L. Vaksman and Ya. S. Soibelman, *The algebra of functions on the quantum group $SU(n+1)$, and odd-dimensional quantum spheres*, Leningrad Math. J. 2 (1991), 1023-1042.
- [We] A. Weinstein, *The local structure of Poisson manifolds*, J. Diff. Geom. 18 (1983), 523-557.
- [Wo1] S. L. Woronowicz, *Twisted $SU(2)$ group: an example of a non-commutative differential calculus*, Publ. RIMS. 23 (1987), 117-181.
- [Wo2] ———, *Compact matrix pseudogroups*, Comm. Math. Phys. 111 (1987), 613-665.